

Note

Perfect One-Factorizations of K_{1332} and K_{6860}

MIDORI KOBAYASHI

Faculty of Economics, Nagasaki University, Nagasaki, 850 Japan

AND

KIYASU-ZEN'ITI

Semiconductor Research Institute, Sendaisi Kawauti, 980 Japan

Communicated by the Managing Editors

Received January 1, 1988

We construct perfect 1-factorizations of K_{1332} and K_{6860} which are newly obtained. © 1989 Academic Press, Inc.

The existence of a perfect 1-factorization of the complete graph K_{2n} for all $n \geq 2$ is conjectured and the problem is settled only for $2n = p + 1$, $2p$ (p is prime), and $2n = 16, 28, 36, 50, 244, 344$ [4-7]. In this note, we construct perfect 1-factorizations of K_{1332} and K_{6860} by applying strong starters to $GF(p^m)$, $p^m \equiv 3 \pmod{4}$.

Let p be a prime number and m be a natural number such that $p^m \equiv 3 \pmod{4}$. We put $q = p^m$, $s = (q - 1)/2$ and $2n = q + 1$. $GF(q)$ denotes the Galois field with q elements and K_{2n} the complete graph with $2n$ vertices.

Let ω be a primitive element of $GF(q)$ and t be an odd integer such that $0 < t < 2s$. We define a starter 1-factor F_0 :

$$F_0 = \{ \{ \omega^{2i}, \omega^{t+2i} \} \mid i = 0, 1, 2, \dots, s-1 \} \cup \{ \{0, \infty\} \}.$$

For any $g \in GF(q)$, we put

$$F_g = F_0 + g = \{ \{ \omega^{2i} + g, \omega^{t+2i} + g \} \mid i = 0, 1, 2, \dots, s-1 \} \cup \{ \{g, \infty\} \}.$$

Then

$$F(\omega^t) = \{ F_g \mid g \in GF(q) \}$$

is a semi-regular 1-factorization of $K_{2n}[2, 3]$.

By suitable selections of the semi-regular 1-factorizations, we construct perfect 1-factorizations.

In case $p = 11$ and $m = 3$, let ω be a primitive element of $GF(11^3)$ with an minimal polynomial $x^3 + x^2 + 5$. Then $F(\omega^7)$ is a perfect 1-factorization of K_{1332} .

In case $p = 19$ and $m = 3$, let ω be a primitive element of $GF(19^3)$ with a minimal polynomial $x^3 + x^2 + 16$. Then $F(\omega^{127})$ is a perfect 1-factorization of K_{6860} .

These perfect 1-factorizations are newly obtained, to the best of the authors' knowledge.

ACKNOWLEDGMENTS

The authors acknowledge Professor G. Nakamura, Dr. J. H. Dinitz, and the referees for their helpful advice.

REFERENCES

1. B. A. ANDERSON, A class of starter induced 1-factorizations, "Graphs and Combinatorics," Lecture Notes in Mathematics Vol. 406, pp. 180–185, Springer-Verlag, New York, 1974.
2. B. A. ANDERSON AND D. MORSE, Some observations on starters, in "Proceedings, 5th S.-E. Conf. Combinatorics, Graph Theory and Computing, Boca Raton, 1974," Congr. Numer. X, pp. 229–235, Utilitas Math., Winnipeg, 1974.
3. J. H. DINITZ, Some perfect Room squares, *J. Combin. Math. Combin. Comput.* **2** (1987), 29–36.
4. E. IHRIG, E. SEAH, AND D. R. STINSON, A perfect one-factorization of K_{50} , *J. of Comb. Math. and Comb. Computing* **1** (1987), 217–219.
5. M. KOBAYASHI, H. AWOKI, Y. NAKAZAKI, AND G. NAKAMURA, A perfect one-factorization of K_{36} , to appear.
6. E. MENDELSON AND A. ROSA, One-factorizations of the complete graph—A survey, *J. Graph Theory* **9** (1985), 43–65.
7. E. SEAH AND D. R. STINSON, A perfect one-factorization for K_{36} , *Discrete Math.*, in press.